



Goat Wellness Care

New Goats:


- Isolate for 14 days. Monitor for appetite, cough, diarrhea, lameness and abscesses.
- Apply Permethrin CDS topically (1/2cc per 50lb) for ticks and lice.
- Labwork:
 - Fresh manure sample: Fecal egg count for gastro-intestinal parasites.
 - Bloodwork: CAE / CL / Johnes
- Trim hooves
- Vaccinate based on vaccine history

All Goats:

- Trim hooves every 6 months
- Fecal egg count every 6 months
- Check FAMACHA for deworming every 4 months
- Vaccinate yearly

Vaccine:	Disease Protected Against:	When to Give:
CDT	Clostridial C & D, Tetanus	All: annually Kids: 6-8 weeks old, booster in 4 weeks Pregnant Does: Fourth month pregnant
Pneumonia	Rhinotracheitis, parainfluenza 3	As directed by your veterinarian.
Anti-Parasitics	Dose	
Ivomec Injectable 1%	1cc per 50 lbs, orally	Roundworms, barber pole, lungworms, nasal bots
Ivomec Drench 0.08%	6ml per 25 lbs orally	
Panacur (Fenbendazole 10%)	5cc per 50 lbs, orally once a day for 3 days	Roundworms, stomach worms, barber pole, tapeworms
Valbazen (albendazole 11%)	4cc per 50 lbs	Cooperia spp., Ostertagia ostertagi and liver flukes
Permethrin	1cc per 100 lb down topline	Ticks and mites

Sub-cutaneous Injection Sites:



Deworm based on FAMACHA score:

FAMACHA®
 Anaemia guide
 Guide sur l'anémie
 Guía de anemia
 مرشد فقر الدم
 ऐनिमिया संबन्धि निर्देश
 貧血症檢測卡

2005

A(1)	B(2)	C(3)	D(4)	E(5)



VIP Equine Services
Amber Bowen DVM, MS, CVA
Alicia Benson DVM, CVA
(707) 235-0230
www.vip-equine.com